

Making our world more productive



# MAPAX<sup>®</sup> – Best for meat and meat products.





Comparison of two different meat packages – one with modified atmosphere packaging and the other one without

Linde's MAPAX® portfolio meets today's food preservation challenges with bespoke gases and mixtures, application expertise and complementary installation, test and safety services.

## Bacteria grow easily on fresh meat

Meat and meat products are particularly susceptible to bacterial growth owing to their high water activity and nutrient content. Meat is sterile to begin with, but when carved up, the surfaces exposed to ambient air provide excellent conditions for the growth of bacteria. Minced meat is naturally even more exposed. For this reason, hygiene and effective temperature control in processing and prepackaging – keeping tools and equipment clean – are vitally important to minimise the contamination of the product with microorganisms.


## Red meat requires oxygen

A special problem arises with red meat such as beef with regard to colour changes caused by the

oxidation of the red pigment. The atmosphere for fresh meat therefore normally contains high concentrations of oxygen (60–80%) in order to retain the red colour by ensuring high oxygen levels in the meat's myoglobin. Highly pigmented meat such as beef thus requires higher oxygen concentrations than low pigmented meat such as pork. With the right mixtures, the practical shelf-life of consumer-packed meat can be extended from 2–4 days to 5–8 days at 4°C.

## The effectiveness of carbon dioxide

Generally speaking, carbon dioxide has a strong inhibiting effect on the growth of bacteria, of which the aerobic genus *Pseudomonas* presents the greatest problem for fresh meat.



Bacteria count over time for meat stored in air and in a modified atmosphere at the same temperature. The meat stored in air enters the period of extremely fast growth, the lag phase, well ahead of the meat stored in the modified atmosphere. This is because the  $\text{CO}_2$  in the modified atmosphere has dissolved into the surface of the meat, reducing its pH value, inhibiting bacterial growth and thus extending the lag phase, until the point when the inhibiting effects become insufficient to control the bacteria.

### Recommended gas mixtures for meat and meat products

Product	Gas mixture	Gas volume Product volume	Typical shelf-life		Storage temp.
			Air	MAP	
Raw red meat	60–80% $\text{O}_2$ + 20–40% $\text{CO}_2$	100–200 ml 100 g meat	2–4 days	5–8 days	2–3 °C
Minced meat	80–100% $\text{O}_2$ + 0–20% $\text{CO}_2$	100–200 ml 100 g meat	<24 h	3–4 days	2–3 °C
Raw light poultry	40–100% $\text{CO}_2$ + 0–60% $\text{N}_2$	100–200 ml 100 g meat	4–7 days	16–21 days	2–3 °C
Raw dark poultry	70% $\text{O}_2$ + 30% $\text{CO}_2$	100–200 ml 100 g meat	3–5 days	7–14 days	2–3 °C
Sausages	20–30% $\text{CO}_2$ + 70–80% $\text{N}_2$	50–100 ml 100 g prod.	2–4 days	2–5 weeks	4–6 °C
Sliced cooked meat	30% $\text{CO}_2$ + 70% $\text{N}_2$	50–100 ml 100 g prod.	2–4 days	2–5 weeks	4–6 °C

## Poultry

Poultry is very susceptible to bacterial spoilage, evaporation loss, off-odour, discolouration and biochemical deterioration. The sterile poultry tissue becomes contaminated during the evisceration process. The practical shelf-life of gas-packed poultry is about 16 to 21 days. The head-space volume should be nearly as large as the product volume. In contrast to red meats, poultry does not undergo irreversible discolouration of the meat's surface in the presence of  $O_2$ . The spoilage of raw poultry is mainly caused by microbial growth, particularly growth of the *Pseudomonas* and *Achromobacter* species. These aerobic spoilage bacteria are very effectively inhibited by  $CO_2$  in modified atmosphere packaging (MAP). Levels of  $CO_2$  in excess of 20% are required to significantly extend the shelf-life of poultry. Package collapse and excessive drip could be a problem for raw poultry, so if higher levels of  $CO_2$  are used, the gas/product ratio should also be increased. Where package collapse is not a problem (e.g.


bulk or master bags), 100%  $CO_2$  is recommended. In both retail and bulk modified atmospheres,  $N_2$  is used as an inert filler gas.

## Meat products have different microflora

Deterioration of meat products is most commonly caused by microbial spoilage. Due to the processing operations, for instance marinating, drying, smoking, fermentation, curing and cooking, the microflora in meat products differ from those in raw meat and the spoilage mechanisms are thereby different. This affects the gas composition used in the package. In order to prevent the products from going off, the concentration of carbon dioxide is usually low (20–50%).



## Microbiological shelf-life of chicken at different atmosphere/temperature combinations



## Linde GmbH

Gases Division, Dr.-Carl-von-Linde-Strasse 6-14, 82049 Pullach, Germany

Phone: +49 89 7446 0, [www.linde-gas.com](http://www.linde-gas.com)

Linde is a company name used by Linde plc and its affiliates. The Linde logo, the Linde word and MAPAX are trademarks or registered trademarks of Linde plc or its affiliates. Copyright © 2016, Linde plc.